

Review Article

Lifestyle changes for prevention of breast cancer

Seyed Hesam Bani Hashemi¹, Samieh Karimi², Hamidreza Mahboobi^{3,4}

¹. Department of surgery, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

². Hormozgan fertility and infertility research center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

³. Infectious and Tropical Diseases Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

⁴. B.A of psychology, Payam Noor University (PNU), Tehran, Iran

Corresponding Author:

Hamidreza Mahboobi, Infectious and Tropical Diseases Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran. Tel: +98.9172121995, Email: hamidrezamahboobi@yahoo.com

Abstract

Breast cancer is the second most common cause of death from cancer among women. Lifestyle changes are shown to be important in the prevention of breast cancer. Diet, physical activity, smoking, alcohol use, and vitamin and mineral use are key factors influencing the risk of breast cancer among women. Because these factors are related to each other, it is difficult to assess their individual roles in breast cancer. Some of these factors are alterable, meaning that women can decrease their risk for breast cancer by changing their behavior. Breast cancer is associated with a high rate of mortality and morbidity among women. Therefore, it is logical to try to find ways to decrease the risk of developing breast cancer. Lifestyle changes seem to be an easy, effective, and economical way to help prevention breast cancer. In women with a confirmed breast cancer diagnosis who are under radiotherapy treatment after undergoing a mastectomy, lifestyle changes are still very important. Some factors, such as smoking cessation and prevention of weight gain, may improve the long-term survival chances of these patients. Therefore, ways to increase women's knowledge about the role of lifestyle changes in the prevention of breast cancer and in the survival of patients with diagnosed breast cancer should be considered and studied.

Keywords: breast cancer, prevention, lifestyle

Bibliographic Information of this article:

[Seyed Hesam Bani Hashemi, Samieh Karimi, Hamidreza Mahboobi. **Lifestyle changes for prevention of breast cancer**. *Electronic Physician*, 2014; 6 (3): 894-905]. (ISSN: 2008-5842). <http://www.ephysician.ir>. Available from: <http://www.ephysician.ir/2014/894-905.pdf>



This work is licensed under a Creative Commons Attribution 3.0 Unported License.

Received: April.29.2014

Revised: 1st Revision: May.17.2014; 2nd Revision: May.20.2014; 3rd Revision: June.09.2014

Accepted: June.10.2014

Published: XXX

1. Introduction

Cancer is a significant cause of mortality and morbidity in the world. Breast cancer is the most common cancer and the second most common cause of death from cancer in women. Breast cancer is an important topic for researchers who are focused on disease prevention. Available screening tests and known risk factors make breast cancer an interesting disease for studies on its prevention. Prevention of breast cancer is possible by changing one's lifestyle. Several important risk factors, which can be altered in the lifestyle of patients, are known to be effective in the prevention of breast cancer.

Many researchers have studied the role of lifestyle change in the prevention of breast cancer. In recent years, several large cohort studies and case control studies with high sample sizes were published. New research has studied the role of lifestyle change in the prevention of breast cancer with more details. It is necessary to review the

recent advances in this field to know where the gaps in knowledge are and to highlight controversies that require more study.

2. Discussion

Lifestyle changes and suitable and on-time screening tests are the most important factors in reducing the rate of breast cancer and in early diagnosis and treatment of breast cancer. Several studies have reported the possible risk factors of breast cancer. Not all of these risk factors are alterable. To prevent breast cancer, it is necessary to know the alterable risk factors and to change them. Studies that have shown important risk factors of breast cancer seem to be adequate and, despite the diversity in the results of studies in different areas of the world, the main risk factors are similar. Although many studies have highlighted the risk factors for breast cancer, few studies have reported strategies for changing these factors. Also, it is unclear whether or not changing these risk factors is enough to decrease the risk of breast cancer. In the other words, studies have mainly focused on identifying the risk factors of breast cancer and recommending altering these behaviors to decrease the risk of breast cancer.

Advantages of alterations of lifestyle are not only limited to women who are just at risk of breast cancer, but also to women who have been diagnosed. Lifestyle change is possible and effective in breast cancer survivors for prevention of recurrence (1). Also, increased survival and improved quality of life have been reported for these women. Fortunately, several recent studies have discussed the role of lifestyle changes in female survivors of breast cancer.

In contrast to studies of healthy women, the results of studies of survivors of breast cancer are more applicable because these studies have often reported the results of their interventions. For example, group-based lifestyle change can be implemented by cancer survivors in order to change the risk factors for poor survival and recurrence (2). The main cause of the fact that stronger reports for the outcomes of lifestyle change are available in women who have survived from breast cancer in comparison to healthy women is that the outcomes can be assessed easier. This is due to the short duration between intervention and outcome. Many risk factors play a role in breast cancer and a large number are related to lifestyle. Therefore, it is impossible to discuss these factors together. Many of these factors interact and may either increase or decrease each other's influence on the risk of breast cancer. Therefore, it seems rational to discuss each factor separately.

2.1. Socioeconomic status

High socioeconomic status is associated with an increased risk of breast cancer (3); several factors play a role in this association. Women with high socioeconomic status make more frequent visits to doctors and have physical examinations, for early diagnosis of breast cancer. Also the lowest number of children increases the risk of breast cancer. The role of socioeconomic status in breast cancer is complex. In some countries, it is shown that women of low socioeconomic status are at a higher risk for diagnosis of breast cancer in the later stages (4, 5). This fact not only confirms inadequate screening in this population, but also means a poorer prognosis in this population at the time of diagnosis. Although women with low socioeconomic status are at a decreased risk of breast cancer, routine screenings for breast cancer are still recommended (5). Women of low socioeconomic status have less access to health care providers in comparison with women with high socioeconomic status in the same area and with the same insurance system (6).

Socioeconomic status changes other factors such as using screening tools, parity, diet, smoking, and alcohol use, which are important factors in the evaluation of the risk of breast cancer in women (3). Therefore, the role of socioeconomic factors in breast cancer is difficult to evaluate independently (7). Individuals with different socioeconomic statuses have some variation in their lifestyles, which changes their risk for breast cancer. Lifestyle changes could be different in women with high and low socioeconomic status. In women with high socioeconomic status, the focus should be on changing dietary habits, smoking cessation, and discontinuation of alcohol consumption. Women with low socioeconomic status may benefit more from regular doctor visits, physical examinations, and screenings for breast cancer.

2.2. Geographical variables

The epidemiological studies in different areas of the world have reported different incidences of breast cancer. The incidence of breast cancer is different even in different areas in the same country. It seems that, in addition to genetic factors, some socioeconomic factors and differences in lifestyle cause variations in the incidence of breast cancer in different areas of the world. North America, Australia/New Zealand, and Western and Northern Europe are reported to be areas with the highest rates of breast cancer. The rate of breast cancer is lower in Asia and Africa (8, 9). Also, there are variations in mortality rates of breast cancer in different geographical areas (10). The differences in the rate of breast cancer and its mortality are correlated with the development of countries. Therefore,

dietary habits, culture, and socioeconomic status are possible explanations for the higher rates in industrial and developed countries. Also, a recent increase in the rate of breast cancer in developing countries confirms this hypothesis.

2.3. Height and weight

The role of body size in the risk of breast cancer is assessed in several studies. Taller women have an increased risk of breast cancer (11, 12). The mechanisms are unclear, but some hormonal factors may be effective. Therefore, tall women should be screened more carefully for breast cancer. In postmenopausal women, high weight is associated with an increase in the risk of breast cancer. This association is more prominent when they do not use hormone therapy. Absence of this association in premenopausal women is in favor of the role of hormones in this association. Obesity is associated with insulin resistance, which seems to be important in assessing the risk of breast cancer. Therefore, hormone therapy in postmenopausal women with a high weight may decrease their risk of breast cancer. Significant weight gain during middle adulthood is associated with an increased risk of breast cancer (13). Some studies have also shown that individuals with a higher weight in premenopausal ages have a lower risk of breast cancer. Studies have controversial results and the mechanisms of this relationship are unclear. Although some studies have shown the beneficial effect of diet and physical activity on weight loss (14), some cohort studies have reported that weight loss and weight gain in women who are diagnosed with breast cancer are associated with limitations in the physical function of women (15). In women with confirmed breast cancer, weight gain is an issue of concern. Several factors such as age, ethnicity, and smoking seem to play a role in this issue (16, 17).

In summary, high height, high weight, and weight gain during middle adulthood are associated with increased risk of breast cancer. However, evidence concerning the role of weight loss in women who are at risk or are diagnosed with breast cancer is inadequate. Therefore, more studies are needed to reach the best intervention in women with high height and high weight.

2.4. Physical activity

Some studies have reported that the risk of breast cancer decreases with increased physical activity (18). Exercise's role in prevention of breast cancer is complicated. The results in premenopausal women are controversial, but, in postmenopausal women, exercise and physical activity decreases the risk for breast cancer by changing the estrogen, insulin, and insulin-like growth factor 1 (IGF-1)(19-21). Women can decrease their risk of breast cancer by engaging in regular exercise. Also, exercise can positively affect other risk factors such as obesity and insulin resistance. It has been shown that post-diagnosis physical activity in women with breast cancer can improve the survival chance of the patients (22, 23). However, it has also been shown that, while the majority of women who are diagnosed with breast cancer will change their diet, they usually do not increase their physical activity (24). It is not clear whether increasing women's knowledge about the role of physical activity in breast cancer will change their level of physical activity or not. A sedentary lifestyle is associated with weight gain and an increased risk of breast cancer. Therefore, physical activity should be recommended to women who are at an increased risk of breast cancer. Women should also be educated about the advantages of physical activity for reducing the risk of breast cancer (25). Physical activity should be recommended to women not only for its role in the prevention of breast cancer, but also for its beneficial roles in managing obesity and insulin resistance. Increasing knowledge may be an effective way to change their physical activity level.

2.5. Smoking

Smoking is an important risk factor for breast cancer. Both active and passive smokers are at an increased risk for breast cancer (26-30). Although other factors of lifestyle and alcohol consumption can alter the relationship between smoking and breast cancer, the evidence is strong enough for a causal relationship between smoking and breast cancer (31). Therefore, smoking is another behavior that can be altered. It is recommended that women stop smoking in order to decrease the risk of breast cancer. Also, smoking increases the chance of other cancers (32) and smoking cessation can decrease the risk of both breast cancer and other cancers.

Women who start smoking at early ages are more susceptible to breast cancer (33). Therefore, teenage girls should be informed of the risk of smoking regarding breast cancer. Parents should also be made aware about this risk. Women who started smoking at early ages should be encouraged to undergo regular screenings for breast cancer. The role of smoking is shown in survivors who have been diagnosed with breast cancer. Women who were smoking at the time of their diagnosis have weaker outcomes and poorer survival (34, 35). However, it still isn't clear that smoking cessation improves the survival of the women who were smokers at the time of the diagnosis of breast cancer. In female smokers who have undergone a partial mastectomy due to breast cancer, the chance for recurrence is about 6.7 times higher than that of women who have never smoked (36). However, more studies are

needed to confirm the role of smoking in the recurrence of breast cancer in women who are undergoing radiation therapy after a partial mastectomy. Smoking and being a passive smoker are some of the most important risk factors for breast cancer and other cancers. It is a risk factor both for breast cancer and its recurrence. Therefore, smoking cessation should be recommended to all women. Enough studies have shown the role of smoking cessation in decreasing the risk of breast cancer, but smoking cessation is difficult and more studies are needed to work on effective strategies of smoking cessation in women who are at risk of breast cancer or who are diagnosed with breast cancer.

2.6. Alcohol

Alcohol consumption is a risk factor for breast cancer; several large studies have shown this association in women (37-39). The risk for breast cancer increases with the amount of alcohol use but is still present with light to moderate alcohol intake (40). The largest cohort study on the relationship between alcohol and breast cancer was done on 105,986 women. Based on the results of this study, small amounts of alcohol use (10 g per day) is associated with a ten percent increase in the risk of breast cancer (41). Similar results were reported in a similar study conducted on 322,000 women. The linear increase in the risk of breast cancer with alcohol consumption is reported in this study (42).

Some studies have reported that folic acid intake attenuates the risk of breast cancer with alcohol consumption (43). More studies are needed about the role of folic acid in prevention of breast cancer in alcoholic women. However, in women who are using alcohol, recommendation to use folic acid is logical. In contrast with smoking, alcohol intake after the diagnosis of breast cancer may not play an important role in survival of the patients (44, 45). However, it is recommended to discontinue alcohol use because the evidence is not enough for a conclusion and because alcohol consumption has other adverse effects on the health (46, 47). Enough evidence is available concerning the risk of breast cancer with alcohol intake. However, more studies are needed to reveal the role of folic acid use and alcohol discontinuation. Therefore, it seems that the recommendation to discontinue alcohol is logical and, in women who are not able to discontinue alcohol, folic acid may be beneficial.

2.7. Fat intake

Fat intake is associated with an increased risk of breast cancer. But there are variations in the results of different studies. Some variables such as amount of fat consumed, reproductive variables, and menopausal age are variables that alter the relationship between fat intake and breast cancer (48-51). Although some cohort studies have reported an increased risk of breast cancer in postmenopausal women with a high fat diet, interventions to reduce fat intake have reported no significant decrease in the risk of breast cancer in healthy postmenopausal women.

The results of studies reporting the role of weight and BMI in the risk of breast cancer are more consistent in comparison to results of studies reporting the role of fat intake in breast cancer. Considering the relationship between fat intake and weight, it seems that the role of weight and BMI is more important than that of fat intake. Although it seems that weight and BMI play a more important role in the risk of breast cancer in comparison to fat intake, there is no difference in the behavior and lifestyle changes that should be recommended to women. Reducing the amount of fat intake reduces the weight and BMI in women, but more studies are needed to show their role in reducing the risk of breast cancer.

2.8. Meat, caffeine, minerals, and vitamins

Some studies have reported an increase in the risk of breast cancer with an increase in the consumption of red meat and caffeine (52-54). The results of studies about caffeine use are inconsistent; some studies reported no association between caffeine consumption and risk of breast cancer (55). It is difficult to assess the role of one specific agent in the risk of breast cancer because of associations between different food materials in different cultures. Breast cancer risk also decreases with an increase in calcium and vitamin D intake (56). This association does not exist with other vitamins and most studies do not support the role of vitamin E and vitamin C in decreasing the risk of breast cancer (57). More studies about the role of vitamins in breast cancer are recommended. There is inadequate evidence about the role of vitamins, minerals, meat, and caffeine in the risk of breast cancer. More research is recommended to increase vitamin intake and restrict meat and caffeine intake.

2.9. Breast feeding

Breastfeeding decreases the risk of breast cancer. Twelve months of breastfeeding is associated with a 4.3 percent decrease in breast cancer risk (58-61). One study in Turkey has reported that shorter durations of breastfeeding can be an important factor that can increase the risk of breast cancer in women. Postmenopausal women who had more than forty-eight months of lactation are at a reduced risk for breast cancer (62). However,

there are other factors, such as parity, that can alter the association between lactation and breast cancer. Even so, it seems that the association is strong enough to recommend breastfeeding in women. Also, breastfeeding has other beneficial effects for both mother and infant.

2.10. Night shift work

Night shift work is known to increase the risk of breast cancer by increasing melatonin levels (63, 64). However, the results of different studies are varied (65). Still more evidence is needed to prove an association between night shift work, melatonin, and breast cancer.

2.11. Stress

While the relationship between stress and other psychological factors and the risk of breast cancer is not strong, it seems that major life events do play a role in the risk of breast cancer (66). It is shown that perceived stress can increase the risk of breast cancer when combined with other risky behaviors such as inadequate physical activity, alcohol intake, smoking, low green tea intake, and high meat and seafood intake (67). Therefore, it is more important for individuals who are in high stress situations to consider nutritional changes and physical activity in order to reduce their risk of breast cancer. In individuals with a breast cancer diagnosis, the level of stress may be higher and the effect of this stress and its control on breast cancer risk may be the subject of future studies in this regard. Also, other psychiatric conditions such as depression may play a role. There are few studies available that assess the role of psychiatric disorders on breast cancer risk.

2.12. NSAID use

The role of NSAIDs in the prevention of breast cancer is not strong, but it seems that NSAID use does decrease the rate of breast cancer (68-70). Some observational studies have reported a decrease in the risk of breast cancer in women who are using NSAIDs, but one randomized controlled trial on the role of aspirin use for prevention of breast cancer reported no significant effects on a 10-year follow up in the study population (71). Research findings do not support the role of NSAIDs in the survival of patients with a definite diagnosis of breast cancer (72), and it seems that the benefits of NSAIDs are limited to individuals who have not been diagnosed with breast cancer. However, more studies are needed to investigate the role of NSAIDs in women who are diagnosed with breast cancer. Although some studies have reported the role of NSAIDs in the prevention of breast cancer, different results were reported in women who were diagnosed with breast cancer. Also, NSAIDs have several side effects affecting the gastrointestinal tract, liver, and kidney. Therefore, using NSAIDs is not recommended for the prevention of breast cancer.

2.13. Digoxin

Several researchers have reported an increase in the risk of breast cancer in patients who are using digoxin (73-75). Women should be informed about this risk before starting digoxin (76). Women who are using digoxin should undergo regular screenings for breast cancer since it is often not possible to discontinue digoxin in patients who are candidates for using it. It seems that digoxin use does not change the risk of recurrence in women with breast cancer who are undergoing treatment (77). Therefore, digoxin can be used in women with breast cancer who need digoxin treatment. They should have regular examinations to watch for possible recurrences. Also, all women should be informed of the risk of digoxin use for breast cancer.

2.14. Importance of education

As noted above, several nutritional changes, physical activity, and alcohol and smoking cessation can reduce the risk of breast cancer in women. Therefore, educating women about these factors and their importance in the risk of breast cancer plays an important role in the prevention of breast cancer (78). Some studies have reported that only giving information about lifestyle changes may not be enough to change the behavior of the female students. A consultancy service may be required in order to change the behaviors of the female students (79). However, it is not clear how useful consultancy service could be. Special strategies, such as group-based education, should be applied to increase the efficacy of women's knowledge in changing their behavior. More studies are needed to show the useful strategies in changing women's behavior. Also, it is important to start this education in early adolescence.

3. Conclusion

As discussed in this review, lifestyle changes are an important factor for prevention of breast cancer. Such changes include reducing weight, adopting a suitable diet, changing unnecessary drugs, etc. Also, in women who are diagnosed and are under treatment for breast cancer, lifestyle changes can reduce the recurrence rate and increase the chance of survival. Women should be informed about the role of lifestyle changes in the prevention of breast cancer and their role in the survival and recurrence rate in patients with breast cancer. More studies are needed to assess the role of education in lifestyle changes, to decrease the incidence and recurrence of breast cancer, and to increase the survival rate.

Acknowledgment

The authors want to thank to Hormozgan University of Medical Sciences for its support in preparing this review.

Conflicts of interest

The authors of this paper declare that they have no conflicts of interest.

Authors' contributions:

All of authors contributed to this project and article equally. All authors read and approved the final manuscript.

References

1. Demark-Wahnefried W, Jones LW, Snyder DC, Sloane RJ, Kimmick GG, Hughes DC, et al. Daughters and Mothers Against Breast Cancer (DAMES): Main outcomes of a randomized controlled trial of weight loss in overweight mothers with breast cancer and their overweight daughters. *Cancer*. 2014. Epub 2014/05/09. <http://dx.doi.org/10.1002/cncr.28761>. PMID: 24804802.
2. Balneaves LG, Van Patten C, Truant TL, Kelly MT, Neil SE, Campbell KL. Breast cancer survivors' perspectives on a weight loss and physical activity lifestyle intervention. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer*. 2014. Epub 2014/03/19. <http://dx.doi.org/10.1007/s00520-014-2185-4>. PMID: 24633590.
3. Pudrovska T, Anishkin A, Shen Y. Early-Life Socioeconomic Status and the Prevalence of Breast Cancer in Later Life. *Research on aging*. 2012;34(3):302-20. Epub 2012/09/01. <http://dx.doi.org/10.1177/0164027511415632>. PMID: 22936839.
4. Wang Q, Li J, Zheng S, Li JY, Pang Y, Huang R, et al. Breast cancer stage at diagnosis and area-based socioeconomic status: a multicenter 10-year retrospective clinical epidemiological study in China. *BMC cancer*. 2012;12:122. Epub 2012/03/30. <http://dx.doi.org/10.1186/1471-2407-12-122>. PMID: 22455370.
5. Farley TA, Flannery JT. Late-stage diagnosis of breast cancer in women of lower socioeconomic status: public health implications. *American journal of public health*. 1989;79(11):1508-12. Epub 1989/11/01. PMID: 2817162.
6. Chang CM, Yin WY, Wei CK, Lin CH, Huang KY, Lin SP, et al. The association of socioeconomic status and access to low-volume service providers in breast cancer. *PloS one*. 2013;8(12):e81801. Epub 2013/12/07. <http://dx.doi.org/10.1371/journal.pone.0081801>. PMID: 24312589.
7. Pruitt SL, Shim MJ, Mullen PD, Vernon SW, Amick BC, 3rd. Association of area socioeconomic status and breast, cervical, and colorectal cancer screening: a systematic review. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2009;18(10):2579-99. Epub 2009/10/10. <http://dx.doi.org/10.1158/1055-9965.EPI-09-0135>. PMID: 19815634.
8. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA: a cancer journal for clinicians*. 2011;61(2):69-90. Epub 2011/02/08. <http://dx.doi.org/10.3322/caac.20107>. PMID: 21296855.
9. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA: a cancer journal for clinicians*. 2005;55(2):74-108. Epub 2005/03/12. PMID: 15761078.
10. Mettlin C. Global breast cancer mortality statistics. *CA: a cancer journal for clinicians*. 1999;49(3):138-44. Epub 1999/08/13. PMID: 10445013.
11. van den Brandt PA, Spiegelman D, Yaun SS, Adami HO, Beeson L, Folsom AR, et al. Pooled analysis of prospective cohort studies on height, weight, and breast cancer risk. *American journal of epidemiology*. 2000;152(6):514-27. Epub 2000/09/21. PMID: 10997541.
12. Lahmann PH, Hoffmann K, Allen N, van Gils CH, Khaw KT, Tehard B, et al. Body size and breast cancer risk: findings from the European Prospective Investigation into Cancer And Nutrition (EPIC). *International journal of cancer Journal international du cancer*. 2004;111(5):762-71. Epub 2004/07/15. <http://dx.doi.org/10.1002/ijc.20315>. PMID: 15252848.
13. Emaus MJ, van Gils CH, Bakker MF, Steins Bisschop CN, Monninkhof EM, Bueno-de-Mesquita HB, et al. Weight change in middle adulthood and breast cancer risk in the EPIC-PANACEA study. *International journal of cancer Journal international du cancer*. 2014. Epub 2014/04/29. <http://dx.doi.org/10.1002/ijc.28926>. PMID: 24771551.
14. Travier N, Fonseca-Nunes A, Javierre C, Guillamo E, Arribas L, Peiro I, et al. Effect of a diet and physical activity intervention on body weight and nutritional patterns in overweight and obese breast cancer survivors. *Med Oncol*. 2014;31(1):783. Epub 2013/12/07. <http://dx.doi.org/10.1007/s12032-013-0783-5>. PMID: 24310809.
15. Young A, Weltzien E, Kwan M, Castillo A, Caan B, Kroenke CH. Pre- to post-diagnosis weight change and associations with physical functional limitations in breast cancer survivors. *Journal of cancer survivorship : research and practice*. 2014. Epub 2014/05/09. <http://dx.doi.org/10.1007/s11764-014-0356-4>. PMID: 24806261.
16. Sedjo RL, Byers T, Ganz PA, Colditz GA, Demark-Wahnefried W, Wolin KY, et al. Weight gain prior to entry into a weight-loss intervention study among overweight and obese breast cancer survivors. *Journal of*

- cancer survivorship : research and practice. 2014. Epub 2014/03/07. <http://dx.doi.org/10.1007/s11764-014-0351-9>. PMID: 24599421.
17. Liu LN, Wen FH, Miaskowski C, Lin YC, Wang JS, Jeng C, et al. Weight change trajectory in women with breast cancer receiving chemotherapy and the effect of different regimens. *Journal of clinical nursing*. 2014. Epub 2014/01/08. <http://dx.doi.org/10.1111/jocn.12521>. PMID: 24393441.
 18. Gao Y, Huang YB, Liu XO, Chen C, Dai HJ, Song FJ, et al. Tea consumption, alcohol drinking and physical activity associations with breast cancer risk among Chinese females: a systematic review and meta-analysis. *Asian Pacific journal of cancer prevention : APJCP*. 2013;14(12):7543-50. Epub 2014/01/28. PMID: 24460331.
 19. Irwin ML, Varma K, Alvarez-Reeves M, Cadmus L, Wiley A, Chung GG, et al. Randomized controlled trial of aerobic exercise on insulin and insulin-like growth factors in breast cancer survivors: the Yale Exercise and Survivorship study. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2009;18(1):306-13. Epub 2009/01/07. <http://dx.doi.org/10.1158/1055-9965.EPI-08-0531>. PMID: 2841479. PMID:19124513.
 20. Fairey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR. Effects of exercise training on fasting insulin, insulin resistance, insulin-like growth factors, and insulin-like growth factor binding proteins in postmenopausal breast cancer survivors: a randomized controlled trial. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2003;12(8):721-7. Epub 2003/08/15. PMID: 12917202.
 21. Ligibel JA, Campbell N, Partridge A, Chen WY, Salinardi T, Chen H, et al. Impact of a mixed strength and endurance exercise intervention on insulin levels in breast cancer survivors. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2008;26(6):907-12. Epub 2008/02/19. <http://dx.doi.org/10.1200/JCO.2007.12.7357>. PMID: 18281663.
 22. Bradshaw PT, Ibrahim JG, Khankari N, Cleveland RJ, Abrahamson PE, Stevens J, et al. Post-diagnosis physical activity and survival after breast cancer diagnosis: the Long Island Breast Cancer Study. *Breast cancer research and treatment*. 2014. Epub 2014/05/03. <http://dx.doi.org/10.1007/s10549-014-2966-y>. PMID: 24789444.
 23. Schmid D, Leitzmann MF. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2014. Epub 2014/03/20. <http://dx.doi.org/10.1093/annonc/mdu012>. PMID: 24644304.
 24. Yaw YH, Shariff ZM, Kandiah M, Weay YH, Saibul N, Sariman S, et al. Diet and physical activity in relation to weight change among breast cancer patients. *Asian Pacific journal of cancer prevention : APJCP*. 2014;15(1):39-44. Epub 2014/02/18. PMID: 24528062.
 25. Catsburg C, Kirsh VA, Soskolne CL, Kreiger N, Bruce E, Ho T, et al. Associations between anthropometric characteristics, physical activity, and breast cancer risk in a Canadian cohort. *Breast cancer research and treatment*. 2014. Epub 2014/05/02. <http://dx.doi.org/10.1007/s10549-014-2973-z>. PMID: 24781974.
 26. Avraham Z, Baron-Epel O, Boker LK. [The relationship between passive smoking, breast cancer risk and n-acetyltransferase 2 (NAT2)]. *Harefuah*. 2014;153(3-4):171-5, 238. Epub 2014/05/06. PMID: 24791559.
 27. Dossus L, Boutron-Ruault MC, Kaaks R, Gram IT, Vilier A, Fervers B, et al. Active and passive cigarette smoking and breast cancer risk: results from the EPIC cohort. *International journal of cancer Journal international du cancer*. 2014;134(8):1871-88. Epub 2014/03/05. <http://dx.doi.org/10.1002/ijc.28508>. PMID: 24590452.
 28. Kawai M, Malone KE, Tang MT, Li CI. Active smoking and the risk of estrogen receptor-positive and triple-negative breast cancer among women ages 20 to 44 years. *Cancer*. 2014;120(7):1026-34. Epub 2014/02/12. <http://dx.doi.org/10.1002/encr.28402>. PMID: 24515648.
 29. Ilic M, Vlajinac H, Marinkovic J. Cigarette smoking and breast cancer: a case-control study in Serbia. *Asian Pacific journal of cancer prevention : APJCP*. 2013;14(11):6643-7. Epub 2014/01/01. PMID: 24377581.
 30. Rosenberg L, Boggs DA, Bethea TN, Wise LA, Adams-Campbell LL, Palmer JR. A prospective study of smoking and breast cancer risk among African-American women. *Cancer causes & control : CCC*. 2013;24(12):2207-15. Epub 2013/10/03. <http://dx.doi.org/10.1007/s10552-013-0298-6>. PMID: 3855285. PMID: 24085586.

31. Nyante SJ, Gierach GL, Dallal CM, Freedman ND, Park Y, Danforth KN, et al. Cigarette smoking and postmenopausal breast cancer risk in a prospective cohort. *British journal of cancer*. 2014;110(9):2339-47. Epub 2014/03/20. <http://dx.doi.org/10.1038/bjc.2014.132>. PMID: 4007228. PMID: 24642621.
32. Land SR, Liu Q, Wickerham DL, Costantino JP, Ganz PA. Cigarette Smoking, Physical Activity, and Alcohol Consumption as Predictors of Cancer Incidence among Women at High Risk of Breast Cancer in the NSABP P-1 Trial. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2014;23(5):823-32. Epub 2014/02/27. <http://dx.doi.org/10.1158/1055-9965.EPI-13-1105-T>. PMID: 4011972. PMID: 24569437.
33. Nishino Y, Minami Y, Kawai M, Fukamachi K, Sato I, Ohuchi N, et al. Cigarette smoking and breast cancer risk in relation to joint estrogen and progesterone receptor status: a case-control study in Japan. *SpringerPlus*. 2014;3:65. Epub 2014/02/12. <http://dx.doi.org/10.1186/2193-1801-3-65>. PMID: 3918095. PMID: 24516791.
34. Berube S, Lemieux J, Moore L, Maunsell E, Brisson J. Smoking at time of diagnosis and breast cancer-specific survival: new findings and systematic review with meta-analysis. *Breast cancer research : BCR*. 2014;16(2):R42. Epub 2014/04/22. <http://dx.doi.org/10.1186/bcr3646>. PMID: 24745601.
35. Pierce JP, Patterson RE, Senger CM, Flatt SW, Caan BJ, Natarajan L, et al. Lifetime cigarette smoking and breast cancer prognosis in the After Breast Cancer Pooling Project. *Journal of the National Cancer Institute*. 2014;106(1):djt359. Epub 2013/12/10. <http://dx.doi.org/10.1093/jnci/djt359>. PMID: 3906992. PMID: 24317179.
36. Bishop JD, Killelea BK, Chagpar AB, Horowitz NR, Lannin DR. Smoking and breast cancer recurrence after breast conservation therapy. *International journal of breast cancer*. 2014;2014:327081. Epub 2014/04/03. <http://dx.doi.org/10.1155/2014/327081>. PMID: 3947862. PMID: 24693439.
37. Scoccianti C, Lauby-Secretan B, Bello PY, Chajes V, Romieu I. Female breast cancer and alcohol consumption: a review of the literature. *American journal of preventive medicine*. 2014;46(3 Suppl 1):S16-25. Epub 2014/02/12. <http://dx.doi.org/10.1016/j.amepre.2013.10.031>. PMID: 24512927.
38. McDonald JA, Goyal A, Terry MB. Alcohol Intake and Breast Cancer Risk: Weighing the Overall Evidence. *Current breast cancer reports*. 2013;5(3). Epub 2013/11/23. <http://dx.doi.org/10.1007/s12609-013-0114-z>. PMID: 3832299. PMID: 24265860.
39. Liu Y, Colditz GA, Rosner B, Berkey CS, Collins LC, Schnitt SJ, et al. Alcohol intake between menarche and first pregnancy: a prospective study of breast cancer risk. *Journal of the National Cancer Institute*. 2013;105(20):1571-8. Epub 2013/08/30. <http://dx.doi.org/10.1093/jnci/djt213>. PMID: 3797023. PMID: 23985142.
40. Park SY, Kolonel LN, Lim U, White KK, Henderson BE, Wilkens LR. Alcohol consumption and breast cancer risk among women from five ethnic groups with light to moderate intakes: the Multiethnic Cohort Study. *International journal of cancer Journal international du cancer*. 2014;134(6):1504-10. Epub 2013/09/17. <http://dx.doi.org/10.1002/ijc.28476>. PMID: 24037751.
41. Chen WY, Rosner B, Hankinson SE, Colditz GA, Willett WC. Moderate alcohol consumption during adult life, drinking patterns, and breast cancer risk. *JAMA : the journal of the American Medical Association*. 2011;306(17):1884-90. Epub 2011/11/03. <http://dx.doi.org/10.1001/jama.2011.1590>. PMID: 3292347. PMID: 22045766.
42. Smith-Warner SA, Spiegelman D, Yaun SS, van den Brandt PA, Folsom AR, Goldbohm RA, et al. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA : the journal of the American Medical Association*. 1998;279(7):535-40. Epub 1998/02/28. PMID: 9480365.
43. Islam T, Ito H, Sueta A, Hosono S, Hirose K, Watanabe M, et al. Alcohol and dietary folate intake and the risk of breast cancer: a case-control study in Japan. *Eur J Cancer Prev*. 2013;22(4):358-66. Epub 2012/11/28. <http://dx.doi.org/10.1097/CEJ.0b013e32835b6a60>. PMID: 23183091.
44. Ali AM, Schmidt MK, Bolla M, Wang Q, Gago-Dominguez M, Esteban Castela J, et al. Alcohol Consumption and Survival after a Breast Cancer Diagnosis: A Literature-based Meta-analysis and Collaborative Analysis of Data for 29,239 Cases. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2014. Epub 2014/03/19. <http://dx.doi.org/10.1158/1055-9965.EPI-13-0901>. PMID: 24636975.
45. Kwan ML, Chen WY, Flatt SW, Weltzien EK, Nechuta SJ, Poole EM, et al. Postdiagnosis alcohol consumption and breast cancer prognosis in the after breast cancer pooling project. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by*

- the American Society of Preventive Oncology. 2013;22(1):32-41. Epub 2012/11/15. <http://dx.doi.org/10.1158/1055-9965.EPI-12-1022>. PMID: 23150063.
46. Newcomb PA, Kampman E, Trentham-Dietz A, Egan KM, Titus LJ, Baron JA, et al. Alcohol consumption before and after breast cancer diagnosis: associations with survival from breast cancer, cardiovascular disease, and other causes. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2013;31(16):1939-46. Epub 2013/04/10. <http://dx.doi.org/10.1200/JCO.2012.46.5765>. PMID: 3661933. PMID: 23569314.
 47. Gou YJ, Xie DX, Yang KH, Liu YL, Zhang JH, Li B, et al. Alcohol Consumption and Breast Cancer Survival: A Meta- analysis of Cohort Studies. *Asian Pacific journal of cancer prevention : APJCP*. 2013;14(8):4785-90. Epub 2013/10/03. PMID: 24083744.
 48. Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *International journal of cancer Journal international du cancer*. 1975;15(4):617-31. Epub 1975/04/15. PMID: 1140864.
 49. Hunter DJ, Spiegelman D, Adami HO, Beeson L, van den Brandt PA, Folsom AR, et al. Cohort studies of fat intake and the risk of breast cancer--a pooled analysis. *The New England journal of medicine*. 1996;334(6):356-61. Epub 1996/02/08. <http://dx.doi.org/10.1056/NEJM199602083340603>. PMID: 8538706.
 50. Cho E, Spiegelman D, Hunter DJ, Chen WY, Stampfer MJ, Colditz GA, et al. Premenopausal fat intake and risk of breast cancer. *Journal of the National Cancer Institute*. 2003;95(14):1079-85. Epub 2003/07/17. PMID: 12865454.
 51. Boyd NF, Stone J, Vogt KN, Connelly BS, Martin LJ, Minkin S. Dietary fat and breast cancer risk revisited: a meta-analysis of the published literature. *British journal of cancer*. 2003;89(9):1672-85. Epub 2003/10/30. <http://dx.doi.org/10.1038/sj.bjc.6601314>. PMID: 2394401. PMID: 14583769.
 52. Taylor EF, Burley VJ, Greenwood DC, Cade JE. Meat consumption and risk of breast cancer in the UK Women's Cohort Study. *British journal of cancer*. 2007;96(7):1139-46. Epub 2007/04/05. <http://dx.doi.org/10.1038/sj.bjc.6603689>. PMID: 2360120. PMID: 17406351.
 53. Ganmaa D, Willett WC, Li TY, Feskanich D, van Dam RM, Lopez-Garcia E, et al. Coffee, tea, caffeine and risk of breast cancer: a 22-year follow-up. *International journal of cancer Journal international du cancer*. 2008;122(9):2071-6. Epub 2008/01/10. <http://dx.doi.org/10.1002/ijc.23336>. PMID: 18183588.
 54. Ishitani K, Lin J, Manson JE, Buring JE, Zhang SM. Caffeine consumption and the risk of breast cancer in a large prospective cohort of women. *Archives of internal medicine*. 2008;168(18):2022-31. Epub 2008/10/15. <http://dx.doi.org/10.1001/archinte.168.18.2022>. PMID: 2574428. PMID: 18852405.
 55. Fagherazzi G, Touillaud MS, Boutron-Ruault MC, Clavel-Chapelon F, Romieu I. No association between coffee, tea or caffeine consumption and breast cancer risk in a prospective cohort study. *Public health nutrition*. 2011;14(7):1315-20. Epub 2011/04/07. <http://dx.doi.org/10.1017/S1368980011000371>. PMID: 21466740.
 56. Shin MH, Holmes MD, Hankinson SE, Wu K, Colditz GA, Willett WC. Intake of dairy products, calcium, and vitamin d and risk of breast cancer. *Journal of the National Cancer Institute*. 2002;94(17):1301-11. Epub 2002/09/05. PMID: 12208895.
 57. Lin J, Cook NR, Albert C, Zaharris E, Gaziano JM, Van Denburgh M, et al. Vitamins C and E and beta carotene supplementation and cancer risk: a randomized controlled trial. *Journal of the National Cancer Institute*. 2009;101(1):14-23. Epub 2009/01/01. <http://dx.doi.org/10.1093/jnci/djn438>. PMID: 2615459. PMID: 19116389.
 58. Stuebe AM, Willett WC, Xue F, Michels KB. Lactation and incidence of premenopausal breast cancer: a longitudinal study. *Archives of internal medicine*. 2009;169(15):1364-71. Epub 2009/08/12. <http://dx.doi.org/10.1001/archinternmed.2009.231>. PMID: 2888088. PMID: 19667298.
 59. Zheng T, Holford TR, Mayne ST, Owens PH, Zhang Y, Zhang B, et al. Lactation and breast cancer risk: a case-control study in Connecticut. *British journal of cancer*. 2001;84(11):1472-6. Epub 2001/06/01. <http://dx.doi.org/10.1054/bjoc.2001.1793>. PMID: 2363665. PMID: 11384096.
 60. Tryggvadottir L, Tulinius H, Eyfjord JE, Sigurvinnson T. Breastfeeding and reduced risk of breast cancer in an Icelandic cohort study. *American journal of epidemiology*. 2001;154(1):37-42. Epub 2001/06/28. PMID: 11427403.
 61. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet*. 2002;360(9328):187-95. Epub 2002/07/23. [http://dx.doi.org/10.1016/S0140-6736\(02\)09454-0](http://dx.doi.org/10.1016/S0140-6736(02)09454-0). PMID: 12133652.

62. Oran B, Celik I, Erman M, Baltali E, Zengin N, Demirkazik F, et al. Analysis of menstrual, reproductive, and life-style factors for breast cancer risk in Turkish women: a case-control study. *Med Oncol*. 2004;21(1):31-40. Epub 2004/03/23. <http://dx.doi.org/10.1385/MO:21:1:31>. PMID: 15034211.
63. Schernhammer ES, Hankinson SE. Urinary melatonin levels and postmenopausal breast cancer risk in the Nurses' Health Study cohort. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2009;18(1):74-9. Epub 2009/01/07. <http://dx.doi.org/10.1158/1055-9965.EPI-08-0637>. PMID: 19124483.
64. Schernhammer ES, Hankinson SE. Urinary melatonin levels and breast cancer risk. *Journal of the National Cancer Institute*. 2005;97(14):1084-7. Epub 2005/07/21. <http://dx.doi.org/10.1093/jnci/dji190>. PMID: 16030307.
65. Sturgeon SR, Doherty A, Reeves KW, Bigelow C, Stanczyk FZ, Ockene JK, et al. Urinary levels of melatonin and risk of postmenopausal breast cancer: women's health initiative observational cohort. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2014;23(4):629-37. Epub 2014/02/11. <http://dx.doi.org/10.1158/1055-9965.EPI-13-1028>. PMID: 24510738.
66. Butow PN, Hiller JE, Price MA, Thackway SV, Krickler A, Tennant CC. Epidemiological evidence for a relationship between life events, coping style, and personality factors in the development of breast cancer. *Journal of psychosomatic research*. 2000;49(3):169-81. Epub 2000/12/09. PMID: 11110988.
67. Wang L, Liao WC, Tsai CJ, Wang LR, Mao IF, Chen CC, et al. The effects of perceived stress and life style leading to breast cancer. *Women & health*. 2013;53(1):20-40. Epub 2013/02/21. <http://dx.doi.org/10.1080/03630242.2012.732680>. PMID: 23421337.
68. Zhao YS, Zhu S, Li XW, Wang F, Hu FL, Li DD, et al. Association between NSAIDs use and breast cancer risk: a systematic review and meta-analysis. *Breast cancer research and treatment*. 2009;117(1):141-50. Epub 2008/11/04. <http://dx.doi.org/10.1007/s10549-008-0228-6>. PMID: 18979210.
69. Agrawal A, Fentiman IS. NSAIDs and breast cancer: a possible prevention and treatment strategy. *International journal of clinical practice*. 2008;62(3):444-9. Epub 2008/01/16. <http://dx.doi.org/10.1111/j.1742-1241.2007.01668.x>. PMID: 18194278.
70. Kwan ML, Habel LA, Slattery ML, Caan B. NSAIDs and breast cancer recurrence in a prospective cohort study. *Cancer causes & control : CCC*. 2007;18(6):613-20. Epub 2007/04/04. <http://dx.doi.org/10.1007/s10552-007-9003-y>. PMID: 17404892.
71. Cook NR, Lee I-M, Gaziano JM, Gordon D, Ridker PM, Manson JE, et al. Low-dose aspirin in the primary prevention of cancer: the Women's Health Study: a randomized controlled trial. *JAMA : the journal of the American Medical Association*. 2005;294(1):47-55.
72. Wernli KJ, Hampton JM, Trentham-Dietz A, Newcomb PA. Use of antidepressants and NSAIDs in relation to mortality in long-term breast cancer survivors. *Pharmacoepidemiology and drug safety*. 2011;20(2):131-7. Epub 2011/01/22. <http://dx.doi.org/10.1002/pds.2064>. PMID: 21254283.
73. Ahern TP, Tamimi RM, Rosner BA, Hankinson SE. Digoxin use and risk of invasive breast cancer: evidence from the Nurses' Health Study and meta-analysis. *Breast cancer research and treatment*. 2014;144(2):427-35. Epub 2014/02/28. <http://dx.doi.org/10.1007/s10549-014-2886-x>. PMID: 24573543.
74. Biggar RJ, Wohlfahrt J, Oudin A, Hjulter T, Melbye M. Digoxin use and the risk of breast cancer in women. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2011;29(16):2165-70. Epub 2011/03/23. <http://dx.doi.org/10.1200/JCO.2010.32.8146>. PMID: 21422417.
75. Ahern TP, Lash TL, Sorensen HT, Pedersen L. Digoxin treatment is associated with an increased incidence of breast cancer: a population-based case-control study. *Breast cancer research : BCR*. 2008;10(6):R102. Epub 2008/12/06. <http://dx.doi.org/10.1186/bcr2205>. PMID: 19055760.
76. Digoxin and breast cancer. *Prescrire international*. 2013;22(137):98. Epub 2013/05/11. PMID: 23662320.
77. Biggar RJ, Andersen EW, Kroman N, Wohlfahrt J, Melbye M. Breast cancer in women using digoxin: tumor characteristics and relapse risk. *Breast cancer research : BCR*. 2013;15(1):R13. Epub 2013/02/21. <http://dx.doi.org/10.1186/bcr3386>. PMID: 23421975.
78. Pieta B, Chmaj-Wierzchowska K, Opala T. Life style and risk of development of breast and ovarian cancer. *Annals of agricultural and environmental medicine : AAEM*. 2012;19(3):379-84. Epub 2012/10/02. PMID: 23020026.

79. Malak AT, Yilmaz D, Tuna A, Gumus AB, Turgay AS. Relations between breast and cervical cancer prevention behaviour of female students at a school of health and their healthy life style. Asian Pacific journal of cancer prevention : APJCP. 2010;11(1):53-6. Epub 2010/07/03. PMID: 20593930.

Electronic Physician

About Electronic Physician

Electronic Physician (ISSN: 2008-5842) is a peer reviewed journal that publishes original papers, reviews, case reports, and other types of contributions in all areas of basic and clinical medical sciences including community health, internal medicine, surgery, nursing, infectious diseases, epidemiology, human genetics and immunology, biochemistry, neurology, psychiatry, and pediatrics.

Indexing and listing:

"Electronic Physician" is indexed or listed in: Islamic World Science Citation Center (ISC), Malaysia Citation Center (MCC), Index Copernicus, EBSCO Host, Directory of Open Access Journals (DOAJ), Genamics Journal Seek, Open-J Gate, Iran Medex Directory of Biomedical Journals, J Gate, Ind Medica, Geneva Foundation for Medical Education and Research (GFMER), University of Chicago, Arkansas University, Johns Hopkins Libraries, University of Miami, University of Arkansas, University of Zurich, Oregon Health & Science University, Directory of Research Journals Indexing (DRJI), Utrecht University Library, John Brown University online library, University Library of Regensburg (Electronic Journals Library), Universidad Veracruzana, Library catalog of the University of Groningen, University library of Saskatchewan, and more.

Manuscript submission:

All of the submissions should be sent to electronicphysician@gmail.com, and c.c. to info@ephysician.ir. Information about the required files, cover letter, preparing the manuscript, writing the structured abstract, conflict of interest form, and other detail is available in the journal website: <http://www.ephysician.ir>

Research ethics and publication ethics:

We have a Zero Tolerance to academic misconduct including "Duplicate Publication", "Salami Publication" and "Plagiarism". All of the manuscripts are under strict investigation regarding any potential unethical conduct. The "Uniform Requirements for Manuscripts (URM) Submitted to Biomedical Journals" is incorporated into our review and publication process. Please visit <http://www.icmje.org> for more detail. You are encouraged to prepare the reference section of the manuscript using the standard introduced by URM. Visit the Committee on Publication Ethics (COPE) at <http://www.publicationethics.org> and World Association of Medical Editors (WAME) at <http://www.wame.org>, Council of Science Editors (CSE) at <http://www.councilscienceeditors.org> and URM at <http://www.icmje.org> to learn about the publication ethics, review policy of journals, Anti plagiarism policy, writing the acknowledgement section of the article, etc. However you can find most of the practical information in our sample article. Please click here to download the sample article.

Key points in preparing a high quality manuscript:

Writing a great abstract for academic papers is very important. Please download our educational article titled "Writing an eye-catching and evocative abstract for a research article: A comprehensive and practical approach" from <http://www.ephysician.ir/2012/520-524.pdf> and use it for your abstract writing.

Key points in preparing a structured abstract for original articles:

You are suggested to prepare your original articles in a standard form. Please download our educational article titled "Writing for academic journals" from <http://www.ephysician.ir/2012/474-476.pdf> and use it for your manuscript preparation.

Review process:

All of the manuscripts go under review. The expected time between manuscript submission and receiving the first round of review results varies depending on the availability of reviewers and other issues but the authors can expect to have the initial result in one month.

Publication fee:

Currently, the Electronic Physician is free for both readers and authors. It clearly means: No subscription fee, No publication fee, No page charges and No submission fee.